arXiv:1302.4478v2 [nucl-th] 1 Mar 2013

# Low-energy limit of the radiative dipole strength in nuclei

Elena Litvinova
*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824-1321, USA*

Nikolay Belov
*Nuclear Physics Department, St. Petersburg State University, 198504 St. Petersburg, Russia*

(Dated: March 4, 2013)

We explain the low-energy anomaly reported in several experimental studies of the radiative dipole strength functions in medium-mass nuclei. These strength functions at very low gamma-energies correspond to the gamma-transitions between very close nuclear excited states in the quasicontinuum. In terms of the thermal mean-field, the low-energy enhancement of the strength functions in highly-excited compound nuclei is explained by nucleonic transitions from the thermally unblocked single-quasiparticle states to the single-(quasi)particle continuum. This result is obtained within the finite-temperature quasiparticle random phase approximation in the coordinate space with exact treatment of the single-particle continuum and exactly eliminated spurious translational mode. The case of radiative dipole strength functions at the nuclear excitation energies typical for the thermal neutron capture is illustrated for $^{94}$Mo and $^{144}$Nd in comparison to available data.

PACS numbers: 21.10.Pc, 21.60.Jz, 25.40.Lw, 27.60.+j

Experimental and theoretical studies of the nuclear low-energy electric dipole response remain among the challenges of the modern nuclear structure physics and attract an increasing interest because of its astrophysical impact. Radiative strength ($\gamma$-strength) at low energies may enhance the neutron capture rates in the r-process of nucleosynthesis [1, 2] with a considerable influence on elemental abundance distributions. One of the key phases of the r-process nucleosynthesis is capture of a thermal neutron with the subsequent $\gamma$-decay of the compound nucleus. The typical neutron energy in the astrophysical plasma is about 100 keV. Therefore, the description of $\gamma$-emission spectra of a compound nucleus with excitation energies of the order of the neutron separation energy is the central problem. Hauser-Feshbach model is a standard tool for calculations of the radiative neutron capture cross sections [3]. Formally, this model includes all possible decay channels via transmission coefficients. In the gamma-decay channel the corresponding coefficient is determined by the radiative strength function which is usually calculated by one of the phenomenological parameterizations [4–6]. However, in more recent works [1, 2, 7] it has been shown that for the most important electric dipole strength these simple models are not sufficient because they do not account for structural details of the strength at the neutron threshold. Sensitivity of the stellar reaction rates to these details emphasizes the importance of their studies within microscopic self-consistent models.

Another key ingredient for the Hauser-Feshbach calculations is the Brink-Axel hypothesis [8] stating that the $\gamma$-strength does not depend on the nuclear excitation energy, in particular, it is the same for excited and non-excited nuclei. Supposedly true for the giant resonances and for the soft modes like pygmy dipole resonance, this hypothesis is, however, violated for the lowest transition energies. For instance, non-zero strength is systematically observed at very low gamma-energies [9]. Radiative strength functions extracted from various measurements [10–14] show an upbend at $E_\gamma \leq 3$ MeV in light nuclei of Fe-Mo mass region. Studies of Ref. [15] have revealed that this phenomenon, occuring in various astrophysical sites, can have a significant impact on their elemental abundances. Phenomenological approaches approximate the low-energy $\gamma$-strength by the tail of the giant dipole resonance with a temperature-dependent width. This is, however, not justified, because the low-energy $\gamma$-strength originates from underlying physics which is completely different from the giant vibrational motion. Modern microscopic theories have excellent tools for computing probabilities of transitions between the nuclear ground state and excited states, but have common problems to describe transitions between excited states. The general many-body techniques like Green function formalism [16, 17], can be applied to $\gamma$-emission and $\gamma$-absorption in excited states of compound type if it is approximated by a semi-statistical model like a finite-temperature mean-field. In such a case, the transitions are described by the finite-temperature version of the random-phase approximation and its extensions. There exist formulations within discrete model spaces [18–20] and models with exact treatment of the single-particle continuum [21–24].

In this Letter, we explain the mechanism for the enhancement of the low-frequency dipole $\gamma$-transitions between the nuclear excited states in the quasicontinuum. We show, for the first time, that this phenomenon can be quantitatively described in terms of a microscopic many-body approach built on the thermal mean-field description of the compound nucleus. Exact treatment of single-particle continuum at finite temperature and exact elimination of the center-of-mass motion are the two essential ingredients for understanding these dipole $\gamma$-transitions with frequencies $E_\gamma \leq$ 3-4 MeV. First of all, we show that the thermal mean field model provides a reasonable approach to the compound nucleus. At the same time, this model is simple enough to allow a straightforward generalization of the many-body methods for the case of a compound nucleus, in terms of finite temperature corresponding to the nuclear excitation energy. The applicability of the finite-temperature mean-field approach to compound nuclei is usually justified by the fact that the product of the canonical distribution and the nuclear level density entering the integrals for thermal averages is a sharp peak approximating the delta-function of the micro-canonical distribution. Indeed, the nuclear level density has an exponential dependence on energy which, together with the canonical distribution, gives a bell-shaped function of energy. But for real medium-mass and heavy nuclei the typical width of this function is estimated by several MeV, within the conditions corresponding to the thermal neutron capture. However, as it is shown below, the dependence of the nuclear excitation energy on the temperature parameter in the thermal mean-field approach is consistent with their phenomenological dependence and with empirical level densities [25]. This fact is used as an alternative justification for employing the finite-temperature mean-field approach for compound nuclei.

The general concept of the finite-temperature mean-field theory [18, 19, 26] is based on the variational principle of maximum entropy minimizing the thermodynamical potential

$$\Omega(\lambda, T) = E - \lambda N - TS, \tag{1}$$

with the Lagrange multipliers $\lambda$ and $T$ determined by the average energy $E$, particle number $N$ and the entropy $S$. These quantities are the thermal averages involving the generalized one-body density operator $\mathcal{R}$:

$$S = -k\mathrm{Tr}(\mathcal{R}\ln\mathcal{R}), \quad E = \mathrm{Tr}(\mathcal{R}\mathcal{H}), \quad N = \mathrm{Tr}(\mathcal{R}\mathcal{N}), \tag{2}$$

where $\mathcal{H}$ is the nuclear Hamiltonian, $\mathcal{N}$ is the particle number operator, and $k$ is Boltzmann constant. Varying the Eq. (1), one can determine the density operator $\mathcal{R}$ with the unity trace:

$$\mathcal{R} = \frac{e^{-(\mathcal{H}-\lambda\mathcal{N})/kT}}{\mathrm{Tr}\Big[e^{-(\mathcal{H}-\lambda\mathcal{N})/kT}\Big]}, \qquad \mathcal{H} = \frac{\delta E[\mathcal{R}]}{\delta\mathcal{R}}. \tag{3}$$



FIG. 1: The dependence of the excitation energy of the thermally excited nuclei on the temperature parameter within the TRMF model (solid curves), in comparison with the parabolic fit (dashed curves).

The form of the nuclear energy density functional $E[\mathcal{R}]$ is a subject of extensive studies, see, for instance, a review [27]. Density functional theory is supposed, in principle, to provide an exact description of many-body problem [28], if the exact density functional is known, however, nuclear physics is still far from a microscopic derivation of such a functional. A covariant density functional theory (CDFT) [29] is one of the most successful examples since it combines a density dependence through a non-linear coupling between the meson fields [30] with pairing correlations based on an effective interaction of finite range. The Lorentz invariance not only allows one to describe the nuclear spin-orbit coupling, which has an essential influence on the underlying shell structure, in a consistent way, but it also puts stringent restrictions on the number of parameters in the corresponding functionals without reducing the quality of the agreement with experimental data. Although the CDFT cannot provide an exact treatment of the full nuclear dynamics, it is a very good starting point for implementation of correlations which are taken in the mean-field only in a static approximation. The CDFT is very convenient for our purpose providing the self-consistent mean-field description where the nucleonic potentials are calculated from the nucleonic densities. Fig. 1 displays the dependence of the excitation energy $E^*$ for selected nuclei on the temperature parameter $T$ obtained within the thermal relativistic mean-field (TRMF) model with NL3 parametrization [31]. The dependence $E^*(T)$ shows features of the back-shifted Fermi gas (FG) distribution $E^*(T) = \delta + aT^2$, therefore we have found reasonable to make the parabolic fits. The values of the level density parameter $a_{RMF}$ determined from the TRMF model are rather close to the empirical values $a_{FG} \approx A/8$ MeV$^{-1}$. Thus, the temperature of an excited compound nucleus can be obtained from its excitation energy by the self-consistent finite-temperature mean-field calculations. In this work, we consider excitation energies $E^*$ around the neutron separation energies $S_n$ of stable medium-mass and heavy nuclei that corresponds, approximately, to the temperatures around and below 1.5 MeV. At such temperatures the mean-field single-particle energies and wave functions remain almost unchanged.

$\gamma$-emission and $\gamma$-absorption of the compound nuclear system are described as an interaction of the nucleus with a sufficiently weak external electromagnetic field $P$ oscillating with some frequency $\omega$. The interaction with such a field causes small amplitude nuclear oscillations around the static equilibrium, so that the total density matrix $\mathcal{R}$ has an oscillating term, in addition to the static thermal mean-field part $\mathcal{R}^0$:

$$\mathcal{R}(t) = \mathcal{R}^0 + \big[\delta\mathcal{R}e^{-i\omega t} + h.c.\big]. \tag{4}$$



FIG. 2: Schematic picture of the possible lowest-energy single-quasiparticle transitions from the thermally unblocked states in an excited compound (left) and from the "frozen" ones in the ground-state nucleus (right).

Variation $\delta\mathcal{R}$ of the density matrix $\mathcal{R}$ in the external field $P$ obeys, in the local approximation, the following integral equation:

$$\delta\mathcal{R}(x;\omega,T) = \delta\mathcal{R}^{(0)}(x;\omega,T) + \\ + \int dx'dx''\mathcal{A}(x,x';\omega,T)F(x',x'')\delta\mathcal{R}(x'';\omega,T), \qquad (5)$$

where $x$ is a multi-index $x = \{\mathbf{r}, s, \tau, \chi\}$ of spatial coordinate $\mathbf{r}$, spin $s$, isospin $\tau$ and component in the quasiparticle space $\chi$. $F(x,x')$ is the effective nucleon-nucleon interaction, $\mathcal{A}(x,x';\omega,T)$ is the two-quasiparticle propagator in the nuclear medium at finite-temperature and

$$\delta\mathcal{R}^{(0)}(x;\omega,T) = \int dx'\mathcal{A}(x,x';\omega,T)P(x'). \qquad (6)$$



FIG. 3: The E1 $\gamma$-strength function for the thermally excited state of $^{94}$Mo at the temperature parameter $T = T_S = T(S_n)$ It is obtained within the TCQRPA, compared to data [11] and to the $\gamma$-strength for the ground state (T=0, dashed curve).



FIG. 4: Same as in Fig. 3, but for $^{144}$Nd, compared to data from Ref. [32].

The propagator $\mathcal{A}(x,x';\omega,T)$ is the key quantity and ideally has to include all the in-medium and surface effects. In the first approximation we calculate it within the thermal continuum quasiparticle random phase approximation (TCQRPA) in terms of the Matsubara temperature Green functions [16, 17]. The full expression of the TCQRPA propagator in the coordinate space is presented in [23]. As the effects of finite temperature and continuum are supposed to give the leading order contribution to the $\gamma$-strength at $|\omega| \to 0$, at this step we use a non-relativistic approach similar to that of Ref. [23]. Thus, in our first illustrative calculations the nucleonic wave functions result from the phenomenological mean field and the effective nucleon-nucleon interaction $F(x,x')$ has the Landau-Migdal ansatz. The propagator consists of the discrete and continuum parts. The discrete part describes transitions between the single-quasiparticle states in the discrete spectrum, and the continuum part describes transitions from the discrete spectrum states to the continuum. Dashed and solid arrows in Fig. 2 show the low-frequency transitions of both kinds, respectively. The effective occupation probability distribution $\tilde{n}_i(E_i,T)$ has much larger diffuseness at $T = T_S = T(S_n)$ than at T=0, being the following product: $\tilde{n}_i(E_i,T) = v_i^2(T)(1 - n_i(E_i,T))$, where $v_i$ are the occupation numbers of Bogoliubov quasiparticles,

$$n_i(E_i,T) = \frac{1}{1+\exp(E_i(T)/kT)}, \qquad (7)$$

and $E_i$ are the eigenvalues of the single-particle Hamiltonian. The radiative dipole strength function is determined by the quantity $\delta\mathcal{R}$ through its convolution with the electromagnetic dipole operator $P_{E1}$:

$$f_{E1}(E_\gamma,T) = -\frac{8e^2}{27(\hbar c)^3}\mathrm{Im}\int dx P^{\dagger}_{E1}(x)\delta\mathcal{R}(x;\omega,T), \qquad (8)$$

$\omega = E_\gamma + i\Delta, \Delta \to 0$. Formally, Eq. (8) corresponds to $\gamma$-absorption, and $\gamma$-emission strength can be calculated for the "final temperature" $T_f = \sqrt{(E^* - \delta - E_\gamma)/a}$ [32]. However, for $E_\gamma \leq$ 3-4 MeV the $\gamma$-absorption and $\gamma$-emission strength functions almost coincide. Their differences will be discussed elsewhere. Figs. 3 and 4 display the radiative dipole strength functions in $^{94}$Mo and $^{144}$Nd calculated within the TCQRPA at finite and zero temperatures. The values of the finite temperature parameter correspond, in each case, to the $S_n$ value and, hence, to the typical compound nucleus formed after the thermal neutron capture. The finite imaginary part $\Delta$ of the energy variable is taken equal to 200 keV and 150 keV, respectively, in accordance to the energy resolution of the experimental data, in order to make a correct comparison. While the dashed curves (T=0, ground state) show at low $\gamma$-energies the tails of the higher-energy transitions between discrete states (blue dashed arrows in Fig. 2) only due to the non-zero value of $\Delta$, the solid curves give in addition the pure thermal continuum strength which remains finite at $\Delta = 0$. The Figs. 3 and 4 illustrate the behavior of the dipole $\gamma$-strength which is typical for nuclei in these two mass regions: the upbend of the $\gamma$-strength appears as a typical feature of the response of medium-mass nuclei in Fe-Mo region while in heavier nuclei like $^{144}$Nd the $\gamma$-strength comes out rather flat at $E_\gamma \to 0$.

Fig. 2 gives a qualitative interpretation for the low-energy enhancement of the $\gamma$-strength. It is clearly seen that transitions from the thermally unblocked states of the single-particle spectrum to the continuum (solid arrow in Fig. 2, left part T>0) form solely the $\gamma$-strength function at very low transition frequency $E_\gamma$. Such type of transitions is not possible in the ground state (T=0) where the lowest-energy transitions to the continuum have much higher energies (solid arrow in Fig. 2, right part). This schematic picture also explains why the upbend of the $\gamma$-strength becomes less pronounced in heavier systems: the temperature parameter corresponding to the neutron threshold $S_n$ is smaller since the threshold itself is typically lower than in lighter systems. Accordingly, the effective thermal occupation probabilities $\tilde{n}(E_i)$ for the single-nucleon states closest to the continuum decrease that, in turn, decreases the transition probabilities.

Summarizing, we give a theoretical interpretation of the low-energy anomaly in the behavior of the radiative dipole strength in medium-mass and heavy nuclei. We have shown that a microscopic approach to nuclear response with coupling to the continuum and exactly eliminated center-of-mass motion, based on the statistical description of the compound nucleus, gives a very good approximation to the low-energy $\gamma$-strength already on the level of the two-quasiparticle configurations. Application to electric dipole response explains the systematic low-energy enhancement of the $\gamma$-strength on the microscopic level. Thus, it is shown that microscopic nuclear many-body theory can be brought to the domain which was previously dominated by phenomenological approaches. The obtained results may have an important consequence for astrophysics, namely for the approaches to r-process nucleosynthesis: those involving Brink-Axel hypothesis may need to be revised.

The authors are indebted to H. Feldmeier, Yu. Ivanov, E. Kolomeitsev, G. Martínez-Pinedo, V. Tselyaev and V. Zelevinsky for enlightening discussions. Support from Helmholtz Alliance EMMI (E.L.) and GSI Summer Student Program 2009 (N.B.) is gratefully acknowledged.

---

[1] S. Goriely and E. Khan, Nucl. Phys. **A706**, 217 (2002).
[2] S. Goriely, E. Khan, and M. Samyn, Nucl. Phys. **A739**, 331 (2004).
[3] J.J. Cowan, F.-K. Thielemann, J.W. Truran, Phys. Rep. **208**, 267 (1991).
[4] S.G. Kadmenskii, V.P. Markushev, V.I. Furman, Yad. Fiz. **37**, 277 (1983), Sov. J. Nucl. Phys. **37**, 165 (1983).
[5] J. Kopecky, M. Uhl, Phys. Rev. C **41**, 1941 (1990).
[6] S.F. Mughabghab, C.L. Dunford, Phys. Lett. B **487**, 155 (2000).
[7] E. Litvinova *et al.* Nucl. Phys. **A823**, 26 (2009).
[8] D. Brink, Ph.D. thesis, Oxford University, 1955; P. Axel, Phys. Rev. **126**, 671 (1962).
[9] A. Schiller, M. Thoennessen, Atomic Data and Nuclear Data Tables **93**, 549 (2007).
[10] A. Voinov *et al.*, Phys. Rev. Lett. 93, 142504.
[11] M. Guttormsen*et al.*, Phys. Rev. C **71**, 044307 (2005).
[12] A. C. Larsen *et al.*, Phys. Rev. C **76**, 044303 (2007).
[13] E. Algin *et al.*, Phys. Rev. C **78**, 054321 (2008).
[14] M. Wiedeking *et al.*, Phys. Rev. Lett. **108**, 162503.
[15] A.C. Larsen, S. Goriely, Phys. Rev. C **82**, 014318 (2010).
[16] T. Matsubara, Prog. Theor. Phys. **14**, 351 (1955).
[17] A.A. Abrikosov, L.P. Gorkov, I.E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics*, Prentice-Hall, 1963.
[18] P. Ring, L.M. Robledo, J.L. Egido, M. Faber, Nucl. Phys. **A419**, 261 (1983).
[19] H.M. Sommermann, Ann. Phys. **151**, 163 (1983).
[20] Y.F. Niu, N. Paar, D. Vretenar, J. Meng, Phys. Lett. B **681**, 315 (2009).
[21] J. Bar-Touv, Phys. Rev. C **32**, 1369 (1985).
[22] V.A. Rodin, M.G. Urin, PEPAN **31**, 975 (2000).
[23] E.V. Litvinova, S.P. Kamerdzhiev, V.I. Tselyaev, Yad. Fiz. **66**, 584 (2003); Phys. Atomic Nuclei **66**, 558 (2003).
[24] E. Khan, N. Van Giai, M. Grasso, Nucl. Phys. **A731**, 311 (2004).
[25] T. Rauscher, Astrophys. J. Suppl. Ser. **147**, 403 (2003).
[26] A.L. Goodman, Nucl. Phys. **A352**, 30 (1981).
[27] M. Bender, P.-H. Heenen, and P.-G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).
[28] W. Kohn and L. J. Sham, Phys. Rev. **137**, A1697 (1965).
[29] T. Gonzales-Llarena, J. L. Egido, G. A. Lalazissis, and P. Ring, Phys. Lett. B **379**, 13 (1996).
[30] J. Boguta and A. R. Bodmer, Nucl. Phys. **A292**, 413 (1977).
[31] G. A. Lalazissis, J. König, and P. Ring, Phys. Rev. **C55**, 540 (1997).
[32] V.A. Plujko, Nucl. Phys. **A649**, 209c (1999).